


Eribulin

Printed from <https://www.cancerquest.org/patients/drug-reference/eribulin> on 04/23/2024



Brand name: HALAVEN®

IUPAC: 11,15:18,21:24,28 Triepoxy-7,9-ethano-12,15-methano-9H,15H-furo[3,2-i]furo[2',3':5,6]pyrano[4,3 b][1,4]dioxacyclopentacosin-5(4H)-one, 2-[(2S)-3-amino-2-hydroxypropyl]hexacosahydro-3 methoxy-26-methyl-20,27-bis(methylene)-, (2R,3R,3aS,7R,8aS,9S,10aR,11S,12R,13aR,13bS,15S,18S,21S,24S,26R,28R,29aS)-, methanesulfonate (salt)
FDA approval: Yes

[Enlace del fabricante](#)

Usage:


Eribulin mesylate (Halaven®) was approved in November 2010 for the treatment of metastatic breast cancer that has been previously treated with at least one taxane and one anthracycline chemotherapy drug. The drug is given via an intravenous (IV) injection.¹

- ¹Chu, E., & DeVita, V. T. (2015). Physicians' cancer chemotherapy drug manual 2015. Burlington, MA: Jones & Bartlett Learning.

Mechanism:

Eribulin looks to cause apoptosis by inducing an irreversible mitotic blockage. Eribulin also works to limit tumor growth by disallowing the creation of microtubules through blocking of spindle formation. Eribulin also helps in the reduction of hypoxic regions by increasing blood flow to the tumor and therefore limiting abnormality and growth.^{1 2}

The diagram below shows the 3D molecular structure of Eribulin.



- ¹Halaven Mechanism of Action for Metastatic Breast Cancer. (2016, November 30). Retrieved June 27, 2017, from <http://www.halaven.com/hcp/metastatic-breast-cancer/MOA>
- ²Halaven Mechanism of Action for Metastatic Breast Cancer. (2016, November 30). Retrieved June 27, 2017, from <http://www.halaven.com/hcp/metastatic-breast-cancer/MOA>

Side effects:

Side effects of this drug may include nausea, stomach problems, constipation, tiredness (fatigue), hair loss, decreased white blood counts (neutropenia) and pain or numbness of the hands and feet (peripheral neuropathy). The drug can lead to an abnormal heart beat. Like any drug that causes decreases in white blood counts, risk of infection may be increased.¹

- ¹Chu, E., & DeVita, V. T. (2015). Physicians' cancer chemotherapy drug manual 2015. Burlington, MA: Jones & Bartlett Learning.

Contraindications:

Eribulin mesylate should not be used by pregnant women or women who could become pregnant. Women using the drug can not breastfeed. Patients with kidney, liver or heart disease should discuss this with their healthcare provider.¹

- ¹Halaven. <http://www.halaven.com/>